



# FibroMeters

- Provide a **score** and a **percentage** of liver fibrosis
- Adapted to the **cause**
  - FibroMeter V (virus B or C or HIV co-infection)
  - FibroMeter A (alcohol)
  - FibroMeter S (metabolic steatosis)
- Improved reliability enabled by an **expert system**




## Hepatitis C


- The expert system allows to affirm or exclude fibrosis or cirrhosis (predictive values at 100%) in a proportion of patients with viral hepatitis

A reliable diagnosis for your patient

## Prediction of significant fibrosis



## Rate of detected cirrhosis by predictive positive values $\geq 95\%$



Data of 1056 chronic hepatitis C from 9 French centres



## Practical information

- ◆ **Easy to prescribe:** ready to use, prescription forms
  - [www.biols.fr](http://www.biols.fr)
  - + 33 (0) 2.41.36.07.19.
  - [contact@biols.fr](mailto:contact@biols.fr)
- ◆ **Results reports** sent by post or available for viewing online.
- ◆ Register to get our **newsletter:** [www.biols.fr](http://www.biols.fr)
- ◆ Also on our **website** : [www.biols.fr](http://www.biols.fr)
  - Scientific information
  - Prescription forms
  - Examples of result reports
  - Also available: personal access codes to prescribe, follow your prescription or view patient results online

Patented research by :



### BioLiveScale

Faculté de Médecine, rue Haute de Reculée 49045 ANGERS Cedex  
Tél. : +33 (0)2.41.36.07.19. / Fax : +33 (0)2.41.36.18.94.  
[www.biols.fr](http://www.biols.fr) / [contact@biols.fr](mailto:contact@biols.fr)

# NEW

Recommended by the French High Health Authority on non-treated chronic hepatitis C without comorbidity

# FibroMeters



## Blood tests for the diagnosis of liver fibrosis

- Virus B or C or HIV co-infection
- Alcohol
- Metabolic steatosis (NAFLD)

