

FibroMeter Ref.: 0904-1410-491

Patient: John MARTIN

Date of birth: 01/03/1970

Date of the blood take: 10/03/2008

Prescriber: Dr. EXAMPLE Physician

University hospital

FR 49100 ANGERS France

Patient's age: 38.0 yrs

Gender: M

### Blood tests results

Markers	Values	Comments
Platelets	315 G/l (*)	(*) Attn, this marker's value is discordant. The expert system has detected 2 markers source of false negative (platelets, prothrombin index).
Prothrombin index	111 % (*)	
ALT	45 UI/l	
AST	97 UI/l	
Urea	4.30 mmol/l	
Hyaluronic acid	64 µg/L	
Alpha 2 macroglobulin	4.27 g/l	

### FibroMeter results


**FibroMeter, score of fibrosis** (scale: 0 to 1):  
is equivalent to the following Metavir stage (scale: 0 to 4):  
(See corresponding meter)

**0.84**  
**F3/4**

Result corrected by the expert system.  
Results should be interpreted with caution  
in accordance with the clinical past-history  
of the patient.

Additional results:


Confirmed presence of liver fibrosis.



Score of fibrosis

**InflaMeter** (scale: 0 to 1):  
is equivalent to the following  
necrotico-inflammatory activity Metavir stage (scale: 0 to 3):  
(See corresponding meter)

**0.39**  
**A1/2**



Activity grade

**Interpretation** - Blood tests allow an exploration of the functional state of fibrosis. As a consequence, because of their construction, they only reflect fibrosis, be it a lesion or morphological (F Metavir type), if the cause is present. Thus, they have diagnostic information. If the cause stops, blood tests decrease faster than the fibrosis lesions. In that case, they do not reflect the lesion but are a predictor of its regression. They, in that case, give prognostic information.

**Expert system** - FibroMeter V score of fibrosis is subject to validation by an "expert system". It detects most of the false negative and false positive results. It then gives a corrected result.

**Precautions** - Blood tests are an additional tool in the diagnosis. As for all medical tests, diagnostic accuracy is not 100%. Results should be confronted to other clinical and para-clinical data by the prescriber